

Thin for quality and health, not spacing

BY MAX BENNETT AND STEPHEN FITZGERALD

Your chainsaw is revved up and you're about to sink the bar into a nearby Douglas fir sapling. But which one? You're surrounded by hundreds of them—some smaller, some larger, all pretty close together, but not uniformly spaced. There are some madrone trees, too, and cedar, and a few spindly pine. You heard somewhere that trees should be 10 feet or 12 feet apart...or was it 15 feet? You can't quite remember. What to do?

Many landowners ask about proper or optimum spacing when thinning trees. We'd suggest not getting too hung up initially on spacing, but instead focus on tree quality and health (vigor). If you leave primarily high-quality trees and take out mostly low-quality trees, it's hard to go wrong. You're likely to have a healthier, more vigorous, more fire-resistant, and more valuable stand in the long run.

Is spacing between trees unimportant? Not at all. Trees need adequate growing space to thrive. However, rigidly adhering to exact spacing (e.g., 14 feet x 14 feet) may lead you to cut trees that should be left and to leave trees that should be cut. This is especially true in natural stands where spacing between trees is highly variable. In plantations where spacing between trees is more regular, using predetermined

spacing or stand density rules makes more sense. Remember that spacing guidelines are just that—they are guidelines and, as such, should remain flexible for leaving the best trees. Also, remember that as trees grow bigger, they progressively need more space, so space trees out wide enough so that they will grow well until the next thinning.

So what makes a high-quality, vigorous tree? Following are some guidelines for "leave" and "cut" trees related to tree quality that you can use to "eyeball" out in the woods. While these points focus on conifer trees, the basic principles also apply to hardwoods.

Characteristics of "leave" trees

- Good live crown ratio (30% or greater)
- Good height growth for species and age
- A-shaped crown (pointy top)
- Abundant foliage with good color
- Good form (straight, without sweeps, crooks, forks, etc.)
- Species is well suited to the site over the long term

Characteristics of "cut" trees

- Poor crown ratio (less than 30%)
- Poor height growth and crown form (flat or rounded top, lopsided)
- Foliage is sparse or off-color

- Poor form or has signs of damage or internal decay
- Species not well suited to the site

Crown ratio

Crown ratio is the percentage of the total height of the tree that is occupied by the live green crown. For example, a tree that is 75 feet tall and has crown on the upper 25 feet of the tree has a live crown ratio of one-third or 33 percent. Crown ratio is important because the bigger the crown (the tree's "food factory"), the better the tree's growth. The rate of growth slows down in most species when the live crown ratio drops below 40 percent. Trees with crown ratios of 30 percent and greater often respond well to release (removal of surrounding competing trees), while trees with crown ratios below 30 percent often respond poorly. These trees may experience thinning shock or sunscald, or grow very slowly. Trees with very small crown ratios may simply die after thinning. This often happens after diameter-limit thinning (removal of all trees above a certain diameter limit) or high grading ("cut the best and leave the rest"). Gradual, light thinnings are recommended for

very dense stands where few, if any, "leave" trees have desirable crown ratios.

Height growth

Height growth can be determined by examining the length of the leader if visible, or on many conifer species (including the true firs, pines, Douglas fir), the length of internodes, which is the distance between branch layers or whorls. Trees with good height growth are preferred as "leave" trees.

Crown shape/form

Pointy-topped or A-shaped crowns indicate rapid height growth. Rounded tops mean height growth has slowed. Good height growth is a sign of good vigor. However, as trees age, height growth slows, and trees tend to develop a round or flat top. Full, symmetrical crowns are preferred over ragged or lopsided crowns. Lopsided crowns develop when a tree is crowded on one or more sides. Trees with lopsided crowns are more vulnerable to blow down or breakage in snow and ice storms.

Crown color/needle density

A dark green color indicates good vigor. Light green or greenish-yellow foliage (chlorotic foliage) is a sign of stress. This may be due to nitrogen deficiency, root disease, bark beetle attacks, or simply moisture stress.

Crown sparseness

A sparse or thinning crown resulting



The tree on the right has a sparse, thinning crown, possibly due to stress, insect attack, or root disease. Trees on the left have full crowns, indicating relatively better health.

from needle loss is another sign of stress. This can often be best determined by comparing a thin or sparse crown with a denser crown on a nearby tree. An abundant crop of undersized cones (distress crop) is sometimes an indication that the tree is severely stressed and likely to die in the near future.

Tree form

The form of the tree trunk (stem) is an important consideration in thinning. All other things being equal, trees with straight trunks and little taper that are free of defect are most desirable as "leave" trees from a timber perspective. However, trees that are very tall for their diameter, especially conifers, tend to be unstable and easily bent by the wind or snow, so some taper is important. Trees that have forked or broken tops, crooked trunks or evidence of rot and defect may be candidates for removal. However, defective trees may also make good wildlife trees, so aiming

for a balance is important. For example, large deformed black oak trees with broken branches or heart rot are some of the best habitat trees around. Unless they threaten to demolish your house or barn if they topple, consider leaving them alone.

Other considerations

These guidelines touch on a few key indicators of tree quality. You should also be on the lookout for evidence of insect and disease problems, some of which may warrant tree removal regardless of other factors. Tree species should also be considered. Many landowners like to leave a mixture of tree species, both for esthetic reasons and because a mixed-species forest is often thought to be less vulnerable to a pest problem that might devastate a single-species stand (monoculture). That's generally true, but make sure that the species is well suited to your site. For example, hot, dry south-facing slopes in the Applegate are good for pine and oak, but aren't so favorable for Douglas fir.

In a nutshell, these guidelines follow the old adage, "cut the sick, lame, and lazy, and leave the better ones." As you thin for quality, you will probably also be reducing stand density to more desirable levels. It may not be too scientific, but forestry, after all, is a science and an art.

Max Bennett

541-776-7371 x221
Forestry/Natural Resources Agent
Oregon State University
Extension Service
Jackson-Josephine Counties
max.bennett@oregonstate.edu

In a nutshell,
... "cut the sick,
lame, and lazy,
and leave the
better ones."

JACKSON COUNTY ANIMAL SHELTER

PETS OF THE MONTH



Chloe - #K0772
Chloe has the cutest markings on her face.



Tristan - #K0415
This big boy looks like a chunky tiger dog.



Emily - #K0854
Emily would love to read poems with you.



Bo - #K1112
This pup has the greatest Spaniel ears.



Hazel - #K0473
Hazel is sure to add a little flavor to your life.



Pippa - #K0385
Pippa is an adorable Rat Terrier girl.

For other informative woodland articles

Visit the online E-Newsletter of SW Oregon Woodland News at <http://extension.oregonstate.edu/sorec/small-woodlands-july-eneews-2012>. Check out the following stories:

- The chain saw circle: Women of the woodlands
- Supreme Court to decide if logging road runoff is pollution
- Environmental groups say it will take 20 years to thin northwest forests
- Forest Capital Partners sells its timber portfolio
- The importance of burls and four other surprises from The Biscuit Fire

fotas 
friends of the animal shelter

adopt ♦ volunteer ♦ donate

WWW.FOTAS.ORG ♦ 541-944-2021

The Jackson County Animal Shelter is located at 5595 S. Pacific Hwy. 99, between Talent & Phoenix. Adoption Hours: 11-4 weekdays, Noon-4 Saturdays.

**LOW-COST
SPAY/NEUTER
COUPONS
AVAILABLE**

www.spayneuter.org
or call 541.858.3325